

The bioenergetic cost of metal resistance and its consequences for reproduction in the harbour ragworm, *Nereis diversicolor*.

Submitted by Christopher Pook to the University of Exeter as a thesis for the degree of *Doctor of Philosophy* in the biological sciences, September 2009.

This thesis is available for Library use on the understanding that it is copyright material and that no quotation from the thesis may be published without proper acknowledgement.

I certify that all material in this thesis which is not my own work has been identified and that no material has previously been submitted and approved for the award of a degree by this or any other University.

.....

Abstract:

The population of harbour ragworms, *Nereis diversicolor*, inhabiting the upper reaches of Restronguet Creek in the Fal Ria, UK, are known to be resistant to the acute, toxic effects of copper, zinc and cadmium. This research aimed to establish whether metabolic and reproductive performance costs were associated with the resistance phenomenon and what the biochemical mechanisms of resistance were, by comparing animals from the Creek with two comparable study sites in the South West of England: Froe Creek and the Teign estuary. There was a significant metabolic cost, measured as a reduction in Scope for Growth, associated with the resistance phenomenon. As no differences in energy intake or uptake were found between resistant and non-resistant animals it was concluded that this cost was covered by demand-side regulation of energetics. A reduction in the amount of biochemical energy reserves in the form of lipids and sugars was also observed in resistant animals suggesting that resource allocation had been shifted away from this endpoint and towards maintenance and activity.

The perturbed metabolism and physiology of resistant *N. diversicolor* is shown to have detrimental consequences for their life-history. Sexually mature, resistant females were significantly smaller than non-resistant ones, indicating that they had either matured at a younger age or grown more slowly. Both total fecundity and mass-specific fecundity were significantly reduced in resistant females, likely as a direct result of the metabolic costs of resistance reducing the resources available to fuel gametogenesis. No differences were found in the energetic reserves stocked within each gamete by resistant and non-resistant animals, suggesting that this trait is heavily selected for.

Synthesis of reduced glutathione [GSH] in resistant animals' tissues appeared to be elevated. Glutathione peroxidase activity was also increased, likely to remediate the effects of Reactive Chemical Species [RCS] that result from the inevitably incomplete binding of intracellular metals. As GSH functions in metal detoxification to covalently binding metals entering resistant animals' cells, preventing their involvement in toxic interactions and their catalysis of the production of RCS, it is proposed that together there two phenomena comprise the fundamental mechanism of resistance to metal toxicity.

Ultimately, this research revealed a hierarchy of health and reproductive performance across the three study populations, with significant associations evident between measured biological endpoints and the degree of metal contamination, illustrating the consequences of anthropogenic pollution for the biology of wild animals.

Table of Contents

	page
List of Figures	9
List of Tables	14
Acknowledgements	15
CHAPTER ONE	17
General introduction to the resistance phenomenon, the Metabolic Cost Hypothesis and the study population.	
1.1 General Introduction.	18
1.1.1 Resistance.	18
1.1.2 Terminology and definitions.	20
1.1.3 Metal resistance.	23
1.1.4 Mechanisms of metal toxicity.	23
1.1.4.1 Mechanisms of metal toxicity: Production of Reactive Chemical Species.	25
1.1.4.2 Mechanisms of metal toxicity: Covalent binding to macromolecules.	27
1.1.5 Mechanisms of metal resistance.	28
1.2 The Metabolic Cost Hypothesis.	31
1.2.1 What are the implications of costs for higher levels of biological organisation?	33
1.2.2 Why is the Metabolic Cost Hypothesis of significance?	36
1.2.3 How do you measure metabolic costs?	38
1.2.4 What previous estimates have been made?	39
1.3 Restronguet Creek.	41
1.3.1 Metal resistance in the <i>N. diversicolor</i> of upper Restronguet Creek.	41

1.3.3	Mechanisms of resistance to metal toxicity.	41
1.3.4	Knowledge gaps.	44
1.4	Aims and hypotheses.	46
1.5	List of abbreviations used.	47

CHAPTER TWO 48

Identification and description of study sites and populations and quantification of resistance to acute copper toxicity.

2.1	General introduction.	50
2.1.1	Description of study sites: Restronguet Creek.	52
2.1.2	Ecology of Restronguet Creek.	54
2.1.3	Description of study sites: Arch Brook in the Teign Estuary.	56
2.1.4	Description of study sites: Froe Creek in the Percuil Estuary.	57
2.1.5	Metal contamination of study sites.	58
2.1.6	Aims of this research.	65
2.2	Methodology for acute toxicity tests.	67
2.3	Results of acute toxicity tests.	71
2.4	Discussion.	73
2.5	List of abbreviations used.	75

CHAPTER THREE 76

Energetic accounting of the metabolic cost associated with metal resistance in *Nereis diversicolor*.

3.1	General introduction to bioenergetics.	8
3.1.1	Bioenergetics in ecotoxicology.	79
3.1.2	The use of simple energy budgets in ecotoxicology.	80
3.1.3	Bioenergetics and the metabolic cost hypothesis.	86

3.1.4	The application of bioenergetics to the study of metal resistance.	88
3.1.5	Aims of this Chapter.	90
3.2	Methodology.	92
3.2.1	Feeding trials.	92
3.2.2	Respirometry.	96
3.2.3	Calculation of Scope for Growth.	98
3.2.4	Quantification of biochemical reserves.	99
3.2.5	Statistical Comparisons.	100
3.3	Results.	101
3.3.1	Physiological measurements.	101
3.3.2	Bioenergetics.	104
3.3.3	Biochemical reserves.	108
3.3.4	Multivariate analysis.	111
3.4	Discussion.	113
3.5	List of abbreviations used.	119

CHAPTER FOUR 120

The reproductive performance cost associated with zinc and copper resistance in *Nereis diversicolor*.

4.1	General introduction.	122
4.1.1	Why study the reproductive biology of <i>N. diversicolor</i> ?	122
4.1.2	Factors affecting reproduction in <i>N. diversicolor</i> .	123
4.1.3	Modelling life-history parameters.	125
4.1.4	Trade-offs between life-history parameters.	126
4.1.5	Aims of this research.	128
4.2	Methodology.	130
4.2.6	Oocyte measurements and fecundity counts.	130
4.2.7	Biochemical composition of oocytes.	133
4.2.8	Energetic assessment of reproductive output.	133
4.2.9	Statistical analysis.	134

4.3	Results.	135
4.3.1	Mature female body size.	135
4.3.2	Oocyte measurements.	136
4.3.3	Incidence of cell death in oocytes.	139
4.3.4	Fecundity of mature females.	140
4.3.5	Oocyte biochemical reserves.	142
4.3.6	Reproductive output.	142
4.3.7	Multivariate analysis.	143
4.4	Discussion.	146
4.5	List of abbreviations used.	153

CHAPTER FIVE 154

Biochemical mechanisms of resistance to metal toxicity in *Nereis diversicolor*.

5.1	General introduction.	156
5.1.1	Metal handling and detoxification.	156
5.1.2	Remediation of metal-induced oxidative stress.	159
5.1.3	Aims of this work.	161
5.2	Methodology.	162
5.2.4	Animals and study sites.	162
5.2.5	Enzyme assays.	162
5.2.6	Glutathione assay.	164
5.2.6.1	Thiol blocking and derivatisation.	165
5.2.6.2	Liquid chromatography.	166
5.2.7	Statistical analysis.	166
5.3	Results.	168
5.3.8	Enzymes.	168
5.3.9	Glutathione method development.	172
5.3.10	Glutathione assay results.	175
5.3.11	Multivariate analysis of biochemistry results.	180
5.4	Discussion.	182

5.5	List of abbreviations used.	187
CHAPTER SIX		187
General discussion.		
6.1	General discussion.	188
6.1.1	Glutathione stoichiometry.	189
6.1.2	Bioenergetic accounting of metal detoxification mechanisms.	192
6.1.3	Effects of resistance upon life history trade-offs.	195
6.1.4	Interpopulation differences between non-resistant populations.	204
6.2	Overall conclusions.	205
6.3	Suggestions for future work.	206
6.4	List of abbreviations used.	208
CHAPTER SEVEN		209
References cited.		
CHAPTER EIGHT		234
Appendix		
8.1	Pook C., Lewis C.N. & Galloway T.S. (2009). The metabolic and fitness costs associated with metal resistance in <i>Nereis diversicolor</i> . Marine Pollution Bulletin 58: 1063-1071	235

List of Figures

<i>figure</i>	<i>page</i>
1.1: Diagram showing various cellular metal transport and handling mechanisms.	30
1.2: General model of resource allocation, redrawn from Sibly and Calow (1990).	35
2.1: Map showing the location of the three populations sampled (Google Maps).	51
2.2: Map of the Fal Ria showing the catchments of its various branches (dotted lines). The locations of historical hard rock mines (filled circles) and smelters (open circles) are indicated.	53
2.3: Silhouette of the lower Fal Ria illustrating the gradient of copper contamination originating at the head of Restronguet Creek.	55
2.4: Satellite Image of Restronguet Creek (Google Maps). The sampling area is shown as a white bar.	55
2.5: Satellite image of the Teign Estuary (Google Maps). The sampling site at Arch Brook is indicated by the white bar.	56
2.6: Satellite image of the Roseland Peninsula and the Percuil Estuary (Google Maps). The sample site at Froe is represented by the white bar.	57
2.7: Map of Restronguet Creek reproduced from Bryan & Gibbs (1983) showing the location of their sample sites (numbered).	58
2.8: Concentrations ($\mu\text{g g}^{-1}$) of copper and zinc in the surface 5cm of sediment in the Fal Ria. From Pirrie et al (2003).	59
2.9: Photograph of the experimental vessels for acute toxicity tests.	69
2.10: Dose-response curves for the acute toxicity of anhydrous copper chloride to three populations of <i>N. diversicolor</i> , generated by probit	72

analysis.

3.1:	Diagram showing the functional links between physiological processes and endpoints and the demographic parameters that ultimately determine fitness. Adapted from Sibly & Calow (1990).	82
3.2:	Photograph of an experimental container containing <i>N. diversicolor</i> . Containers were covered with glass lids to prevent the animals escaping and paper towels to minimise ambient light levels.	93
3.3:	Photograph showing a polychaete in its container. The large green object is an algal pellet, the small green-brown blobs are faecal pellets and the yellow-grey strands are mucus. During collection of faecal material such strands- which were produced by many animals in abundance- were carefully separated from the faecal material.	95
3.4:	Photograph showing a polychaete restrained within its refuge by pieces of 2mm nylon mesh tied across each opening for the measurement of standard metabolic rates.	97
3.5:	Boxplots of data for feeding rate and absorption efficiency measurements for three populations of <i>N. diversicolor</i> .	102
3.6:	Boxplots of data for resting, unfed oxygen consumption for three populations of <i>N. diversicolor</i> .	103
3.7:	Boxplots of data for energy intaken and uptaken for the three study populations of <i>N. diversicolor</i> .	105
3.8:	Boxplots of data for standard metabolic rate and Scope for Growth for three populations of <i>N. diversicolor</i> .	107
3.9:	Boxplots of data for lipid content of three populations of <i>N. diversicolor</i> .	108
3.10:	Boxplots of data for sugar content for the three study populations of <i>N. diversicolor</i> .	109
3.11:	Boxplots of data for protein content for three populations of <i>N. diversicolor</i> .	110
3.12:	Multidimensional Scaling plot of the independent physiological energetics data.	112

4.1:	Image showing oocytes in different stages of maturation. Oocytes labelled “GVBD” show symptoms of germinal vesicle breakdown.	131
4.2:	Image of <i>N. diversicolor</i> oocytes showing predominantly healthy examples. Cells exhibiting symptoms of cell death (swelling and lifting of the plasma membrane) have been labelled (N).	132
4.3:	Wet weights of mature females from three populations of <i>N. diversicolor</i> .	136
4.4:	Diameters of oocytes collected from mature females sampled from the three study populations.	138
4.5:	Proportion of dying oocytes observed in <i>N. diversicolor</i> from the three study populations.	139
4.6:	Results of the total fecundity and mass-specific, effective fecundity measurements made upon mature females from three populations of <i>N. diversicolor</i> .	141
4.7:	Boxplots showing the Specific, Effective Reproductive Output of female <i>N. diversicolor</i> from the three study populations.	143
4.8:	Multidimensional Scaling plot of the independent reproductive parameters measured here.	145
5.1:	Boxplots of data for glutathione peroxidase activity in the tissues of <i>N. diversicolor</i> sampled from the three study sites.	168
5.2:	Boxplots of data for glutathione reductase activity in the tissues of <i>N. diversicolor</i> sampled from the three study sites.	169
5.3:	Boxplots of data for glutathione transferase activity in the tissues of animals samples from three populations of <i>N. diversicolor</i> .	170
5.4:	Boxplots of data for superoxide dismutase assay in the tissues of animals sampled from three populations of <i>N. diversicolor</i> .	171
5.5:	Standard curves of reduced [GSH] and oxidised [GSSG] glutathione,	172

	quantified as the concentration of reduced glutathione equivalents.	
5.6:	HPLC chromatograms. [A] shows six overlaid chromatograms of glutathione standards of 1.25, 2.5, 5, 10, 20 and 40 μ M concentration of reduced glutathione. [B] shows an experimental assay in <i>N. diversicolor</i> homogenate for total glutathione.	173
5.7:	Boxplots showing the total glutathione content of <i>N. diversicolor</i> sampled from the three study populations.	174
5.8:	Boxplots showing the bound, oxidised glutathione [GSX] content of <i>N. diversicolor</i> sampled from the three study populations.	175
5.9:	Boxplots of data for free, oxidised glutathione [GSSG] concentration for three populations of <i>N. diversicolor</i> .	176
5.10:	Boxplots of data for reduced glutathione concentrations [GSH] in the tissues of <i>N. diversicolor</i> sampled from the three study sites.	177
5.11:	Boxplots of data for the glutathione ratio of the tissues of animals sampled from three populations of <i>N. diversicolor</i> .	178
5.12:	Multidimensional Scaling plot of the independent variables measured in this research (glutathione peroxidase, glutathione reductase, glutathione transferase and superoxide dismutase specific activities and the concentrations of bound, oxidised glutathione, oxidised glutathione homodimer and reduced glutathione.	180
6.1:	Graph of glutathione reductase activity against tissue concentrations of oxidised glutathione for the three study populations.	190
6.2:	Graph of wet weights at maturity against fecundities of individual females from the three study populations.	196
6.3:	Diagram showing the trade-off between growth rate and mortality, redrawn from Sibly and Calow (1989).	198
6.4:	Two models of reproductive value at birth for various levels of non-breeding mortality in <i>N. diversicolor</i> reproducing at ages of one to four years.	202

List of Tables

<i>table</i>	<i>page</i>
2.1: Concentrations ($\mu\text{g g}^{-1}$) of different metals in the sediments at the three experimental sites. “R” – Restronguet Creek, “F” – Froe Creek, “T” – Teign Estuary.	62
2.2: List of reported acute toxicities of metals, as 96 hour median LC50s, to <i>N. diversicolor</i> .	64
2.3: Exposure concentrations of anhydrous copper (II) chloride for acute toxicity tests for ostensibly resistant and non-resistant populations of <i>N. diversicolor</i> , in $\mu\text{g l}^{-1}$.	69
2.4: Dissolved copper LC50s and confidence intervals in $\mu\text{g l}^{-1}$, calculated by probit analysis.	71
4.1: Summary of mean ($\pm\text{SD}$) protein, sugar and lipid contents of individual oocytes collected from sexually mature females sampled from the three study populations of <i>N. diversicolor</i> .	142